

5 METHODS AND COMPOSITIONS RELATING TO CD39-LIKE POLYPEPTIDES  
AND NUCLEIC ACIDS

Ans. A1

1. FIELD OF THE INVENTION

The present invention provides novel polynucleotides and proteins encoded by such polynucleotides, along with therapeutic, diagnostic and research utilities for these  
10 polynucleotides and proteins.

2. BACKGROUND OF THE INVENTION

Technology aimed at the discovery of protein factors (including e.g., cytokines, such as lymphokines, interferons, CSFs and interleukins) has matured rapidly over the past  
15 decade. The now routine hybridization cloning and expression cloning techniques clone novel polynucleotides "directly" in the sense that they rely on information directly related to the discovered protein (i.e., partial DNA/amino acid sequence of the protein in the case of hybridization cloning; activity of the protein in the case of expression cloning). In  
20 addition, more recently, "indirect" cloning techniques have been utilized, such as signal sequence cloning, which isolates DNA sequences based on the presence of a now well-recognized secretory leader sequence motif, as well as various PCR-based or low stringency hybridization cloning techniques, coupled in certain instances with database  
25 searching, have advanced the state of the art by making available large numbers of DNA/amino acid sequences for proteins that are known to have biological activity by virtue of their secreted nature in the case of leader sequence cloning, by virtue of the cell or tissue source in the case of PCR-based techniques and/or by virtue of their sequence  
30 similarity via database searches. It is to these proteins and the polynucleotides encoding them that the present invention is directed.

### 3. SUMMARY OF THE INVENTION

The compositions of the present invention include novel isolated polypeptides, in particular, novel CD-39-like polypeptides, isolated polynucleotides encoding such  
5 polypeptides, including recombinant DNA molecules, cloned genes or degenerate variants thereof, especially naturally occurring variants such as allelic variants, and antibodies that specifically recognize one or more epitopes present on such polypeptides.

The compositions of the present invention additionally  
10 include vectors, including expression vectors, containing the polynucleotides of the invention, cells genetically engineered to contain such polynucleotides and cells genetically engineered to express such polynucleotides.

The isolated polynucleotides of the invention include, but are not limited to, a polynucleotide encoding a  
15 polypeptide comprising the amino acid sequence of SEQ ID NO:2 (also referred to herein as "CD39L2"); or a polynucleotide encoding a polypeptide comprising amino acid residues 72-93, 147-162, 191-211 OR 217-238 of SEQ ID NO:2.

In selected embodiments, such isolated polynucleotides of the invention represents a polynucleotide comprising the  
20 nucleotide sequence of SEQ ID NO:1; or a polynucleotide comprising nucleotides 232-1599, 445-510, 670-717, 802-864 or 880-945 of the nucleotide sequence of SEQ ID NO:1.

The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes to the complement of the nucleotide sequence of  
25 SEQ ID NO:1 under highly stringent hybridization conditions; a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NO:1 under moderately stringent hybridization conditions; a polynucleotide which is an allelic variant of any polynucleotide recited above; a polynucleotide which encodes a species homologue of any of  
30 the proteins recited above; or a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptide of SEQ ID NO:2.

Patented 1994

The isolated polynucleotides of the invention still further include, but are not limited to, a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:4 (also referred to herein as "CD39L3"); or a  
5 polynucleotide encoding a polypeptide comprising amino acid residues 55-76, 132-150, 177-199 or 213-234 of SEQ ID NO:4.

In selected embodiments, such isolated polynucleotides of the invention represents a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3; or a polynucleotide comprising nucleotides 83-1669, 245-310, 476-532, 611-679 or  
10 719-784 of the nucleotide sequence of SEQ ID NO:3.

The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NO:3 under highly stringent hybridization conditions; a polynucleotide that hybridizes to the complement of the  
15 nucleotide sequence of SEQ ID NO:3 under moderately stringent hybridization conditions; a polynucleotide which is an allelic variant of any polynucleotide recited above; a polynucleotide which encodes a species homologue of any of the proteins recited above; or a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of  
20 the polypeptide of SEQ ID NO:4.

The isolated polynucleotides of the invention still further include, but are not limited to, a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:6 (also referred to herein as "CD39L4"); or a  
25 polynucleotide encoding a polypeptide comprising amino acid residues 47-68, 123-138, 167-187 or 193-214 of SEQ ID NO:6; or a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:9 (also referred to herein as dCD39L4"); or a polynucleotide encoding amino acid residues 77-98, 153-167, 197-217 or 223-242 of SEQ ID NO:9.

In one embodiment, such isolated polynucleotides of the  
30 invention represents a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5; or a polynucleotide comprising nucleotides 247-1530, 385-450, 613-660, 745-807 or 823-888 of the nucleotide sequence of SEQ ID NO:5.

100-20-243000

The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NO:5 under highly stringent hybridization conditions; 5 a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NO:5 under moderately stringent hybridization conditions; a polynucleotide which is an allelic variant of any polynucleotide recited above; a polynucleotide which encodes a species homologue of any of the proteins recited above; or a polynucleotide that encodes 10 a polypeptide comprising a specific domain or truncation of the polypeptide of SEQ ID NO:6.

The isolated polynucleotides of the invention further include, but are not limited to, a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:8 (also referred to herein as "mCD39L4" or "mNTPase"); or a 15 polynucleotide encoding amino acid residues 46-67, 122-140, 166-187 or 194-213 of SEQ ID NO:8.

In selected embodiments, such isolated polynucleotides of the invention represents a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7; or a polynucleotide comprising nucleotides 205-1599, 340-395, 568-624, 700-765 or 20 784-843 of the nucleotide sequence of SEQ ID NO:7.

The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NO:7 under highly stringent hybridization conditions; a polynucleotide that hybridizes to the complement of the 25 nucleotide sequence of SEQ ID NO:7 under moderately stringent hybridization conditions; a polynucleotide which is an allelic variant of any polynucleotide recited above; a polynucleotide which encodes a species homologue of any of the proteins recited above; or a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of 30 the polypeptide of SEQ ID NO:8.

The polynucleotides of the invention additionally include the complement of any of the polynucleotides recited above.





invention can further be utilized as anti-thrombotic agents, anti-tissue graft rejection agents, and/or as part of methods for regulating neurotransmission by ATP in smooth muscle, peripheral ganglia or the brain.

5 The methods of the present invention further relate to methods for detecting the presence of the polynucleotides or polypeptides of the invention in a sample. Such methods can, for example, be utilized as part of prognostic and diagnostic evaluation of disorders as recited above and for the identification of subjects exhibiting a predisposition to  
10 such conditions. Furthermore, the invention provides methods for evaluating the efficacy of drugs, and monitoring the progress of patients, involved in clinical trials for the treatment of disorders as recited above.

The invention also provides methods for the identification of compounds that modulate the expression of  
15 the polynucleotides and/or polypeptides of the invention. Such methods can be utilized, for example, for the identification of compounds that can ameliorate symptoms of disorders as recited above. Such methods can include, but are not limited to, assays for identifying compounds and other substances that interact with (e.g., bind to) the  
20 polypeptides of the invention.

The methods of the invention also include methods for the treatment of disorders as recited above which may involve the administration of such compounds to individuals exhibiting symptoms or tendencies related to disorders as recited above. In addition, the invention encompasses  
25 methods for treating diseases or disorders as recited above by administering compounds and other substances that modulate the overall activity of the target gene products. Compounds and other substances can effect such modulation either on the level of target gene expression or target protein activity.

30

#### 4. BRIEF DESCRIPTION OF THE FIGURES

Figure 1. Top line: Nucleotide sequence of SEQ ID NO:7, referred to herein as mNTPase or mCD39L4; bottom line:

amino acid sequence of SEQ ID NO:8, referred to herein as mNTPase or mCD39L4.

Figure 2. Amino acid alignments of the full mNTPase (mCD39L4) amino acid sequence (SEQ ID NO:8) and the most closely related other NTPase proteins: garden pea NTPase (SEQ ID NO:10), potato apyrase (SEQ ID NO:11), yeast GDPase (SEQ ID NO:12). Identical residues are indicated by dark background (white letters), while conserved residues are indicated by gray background. Alignments were made with pileup and boxshade from the Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison WI.

Figure 3. Alignment of 12 members of the NTPase (or CD39-like) gene family indicating the conserved apyrase regions I-IV. CD39=human (from Accession No. S73813; SEQ ID NO:13), ratCD39=rat (from Accession No. gi11754710; SEQ ID NO:14), CD39L1=human (Accession No. U91510; SEQ ID NO:15), ChickATPase=chicken (from Accession No. U74467; SEQ ID NO:16), peaNTPase=garden pea (from Accession No. P52914; SEQ ID NO:10), potRROP1=potato RROP1 gene (from Accession No. gi11381633; SEQ ID NO:11), yGDA1+y71KD=yeast genes (from Accession Nos. sp1P32621 + sp1P40009; SEQ ID NO:12), hCD39L2=CD39L2, celegans=C. Elegans gene (from Accession No. gi11086594; SEQ ID NO:17). Identical residues are indicated by dark background (white letters), conserved residues are indicated by gray background. Alignments were made with pileup and boxshade from the Wisconsin Package 9.0, Genetics Computer Group (GCG), Madison, WI. Conserved portions of ACRs I-IV are boxed.

Figure 4. Top line: Nucleotide sequence of SEQ ID NO:1, referred to herein as CD39L2; bottom line: amino acid sequence of SEQ ID NO:2, referred to herein as CD39L2.

Figure 5. Comparison of the hydrophobicity predictions for the amino acid sequences of members of the human CD39-like gene family. Predictions were made using the Topred-II



1.1 program (Claros, M.G. & Von Hejine, G., 1994, Comput. Appl. Biosci. 10:685-686; putative setting=0.5; certain setting=1.0).

5        Figure 6. Top line: Nucleotide sequence of SEQ ID NO:3, referred to herein as CD39L3; bottom line: amino acid sequence of SEQ ID NO:4, referred to herein as CD39L3.

Figure 7. Top line: Nucleotide sequence of SEQ ID NO:5, referred to herein as CD39L4; bottom line: amino acid  
10 sequence of SEQ ID NO:6, referred to herein as CD39L4.

Figure 8. Amino acid alignments of the full-length protein sequences for human members of the CD39-like gene family. CD39 (from Accession No. S73813; SEQ ID NO:13), CD39L1 (from Accession No. U91510; SEQ ID NO:15), CD39L2 (it  
15 is noted that the CD39L2 polypeptide illustrated here depicts a derived amino acid sequence that is encoded from the ATG codon beginning at nucleotide 148 (see FIG. 4) and, therefore, includes an additional 28 amino acid residues N-terminal to those depicted in FIG. 4; this form of CD39L2 is also intended to be included within the scope of the present  
20 invention), CD39L3, CD39L4.

Identical residues are indicated by a black background (white letters), and conserved residues are indicated by a gray background. Spaces in the sequences are indicated by a dot. Apyrase regions (ACRs) are indicated by arrows, with conserved portions of ACRs I-IV are highlighted by the boxed  
25 sections. Alignments were made using pileup and boxshade from the Wisconsin Package Version 9.0 Genetics Computer Group (GCG), Madison, WI.

Figure 9. Amino acid sequence of dCD39L4 ("dNTPase"; SEQ ID NO:9) and alignment of the amino acid sequence with the  
30 most closely related members of the CD39-like gene family. peaGDP, garden pea NTPase (from Accession No. P52194; SEQ ID NO:10); ptoapyrase, potato RROP1 gene (from Accession No. gill1381633; SEQ ID NO:11); CD39L2; CD39L4, and yGDPase, yeast

yGDA1 gene (from Accession No. sp1P32621; SEQ ID NO:12).  
Apyrase regions (ACRs) are indicated by arrows, with  
conserved portions of ACRs I-IV are highlighted by the boxed  
sections.

5

## 5. DETAILED DESCRIPTION OF THE INVENTION

### 5.1. Definitions

10 The term "nucleotide sequence" refers to a heteropolymer  
of nucleotides or the sequence of these nucleotides. The  
terms "nucleic acid" and "polynucleotide" are also used  
interchangeably herein to refer to a heteropolymer of  
nucleotides. Generally, nucleic acid segments provided by  
this invention may be assembled from fragments of the genome  
and short oligonucleotide linkers, or from a series of  
oligonucleotides, or from individual nucleotides, to provide  
15 a synthetic nucleic acid which is capable of being expressed  
in a recombinant transcriptional unit comprising regulatory  
elements derived from a microbial or viral operon, or a  
eukaryotic gene. In alternate embodiments, a nucleotide  
sequence, polynucleotide or nucleic acid can correspond to a  
genomic sequence (e.g., can contain intron as well as exon  
20 sequence) or cDNA sequences (that is, contains no intron  
sequence).

The terms "oligonucleotide fragment" or a  
"polynucleotide fragment", "portion," or "segment" is a  
stretch of polypeptide nucleotide residues which is long  
enough to use in polymerase chain reaction (PCR) or various  
25 hybridization procedures to identify or amplify identical or  
related parts of mRNA or DNA molecules.

The terms "oligonucleotides" or "nucleic acid probes"  
are prepared based on the polynucleotide sequences provided  
in the present invention. Oligonucleotides comprise portions  
of such a polynucleotide sequence having at least about 15  
30 nucleotides and usually at least about 20 nucleotides.  
Nucleic acid probes comprise portions of such a  
polynucleotide sequence having fewer nucleotides than about 6



is derived from recombinant (e.g., microbial or mammalian) expression systems. "Microbial" refers to recombinant polypeptides or proteins made in bacterial or fungal (e.g., yeast) expression systems. As a product, "recombinant microbial" defines a polypeptide or protein essentially free of native endogenous substances and unaccompanied by associated native glycosylation. Polypeptides or proteins expressed in most bacterial cultures, e.g., *E. coli*, will be free of glycosylation modifications; polypeptides or proteins expressed in yeast will have a glycosylation pattern in general different from those expressed in mammalian cells.

The term "recombinant expression vehicle or vector" refers to a plasmid or phage or virus or vector, for expressing a polypeptide from a DNA (RNA) sequence. An expression vehicle can comprise a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription initiation and termination sequences. Structural units intended for use in yeast or eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport sequence, it may include an N-terminal methionine residue. This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

The term "recombinant expression system" means host cells which have stably integrated a recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit extrachromosomally. Recombinant expression systems as defined herein will express heterologous polypeptides or proteins upon induction of the regulatory elements linked to the DNA segment or synthetic gene to be expressed. This term also means host cells which have stably integrated a recombinant genetic element or elements having a regulatory role in gene expression, for



100-20-245000

the ability of the polypeptide to exhibit an immunological activity. A polypeptide exhibits an "immunological activity" when antibodies can be generated that are directed against the polypeptide.

5 The term "naturally occurring polypeptide" refers to polypeptides produced by cells that have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including, but not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and  
10 acylation.

The term "mature" refers to a polypeptide that has been postranslationally modified or that corresponds in primary amino acid sequence to a polypeptide that has been postranslationally modified. A mature polypeptide includes, but is not limited to, a polypeptide which comprises a  
15 primary amino acid sequence that has been processed from a "pre-," "pro-," or "pre-pro" amino acid sequence; a polypeptide which comprises a primary amino acid sequence corresponding to that of a polypeptide that has been processed from a "pre-," "pro-," or "pre-pro" amino acid sequence; a polypeptide that has been post-translationally  
20 modified via such modifications as, for example, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

The term "derivative" refers to polypeptides chemically modified by such techniques as ubiquitination, labeling (e.g., with radionuclides or various enzymes), pegylation  
25 (derivatization with polyethylene glycol) and insertion or substitution by chemical synthesis of amino acids such as ornithine, which do not normally occur in human proteins.

The term "recombinant variant" refers to any polypeptide differing from naturally occurring polypeptides by amino acid insertions, deletions, and substitutions, created using  
30 recombinant DNA techniques. Guidance in determining which amino acid residues may be replaced, added or deleted without abolishing activities of interest, such as cellular trafficking, may be found by comparing the sequence of the

FOUO 20-CH25000

particular polypeptide with that of homologous peptides and minimizing the number of amino acid sequence changes made in regions of high homology.

Preferably, amino acid "substitutions" are the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, i.e., conservative amino acid replacements. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

"Insertions" or "deletions" are typically in the range of about 1 to 5 amino acids. The variation allowed may be experimentally determined by systematically making insertions, deletions, or substitutions of amino acids in a polypeptide molecule using recombinant DNA techniques and assaying the resulting recombinant variants for activity.

Alternatively, where alteration of function is desired, insertions, deletions or non-conservative alterations can be engineered to produce altered polypeptides. Such alterations can, for example, alter one or more of the biological functions or biochemical characteristics of the polypeptides of the invention. For example, such alterations may change polypeptide characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover rate. Further, such alterations can be selected so as to generate polypeptides that are better suited for expression, scale up and the like in the host cells chosen for expression. For example, cysteine residues can be deleted or substituted with another amino acid residue in order to eliminate disulfide bridges.

U.S. Patent

As used herein, "substantially equivalent" can refer both to nucleotide and amino acid sequences, for example a mutant sequence, that varies from a reference sequence by one or more substitutions, deletions, or additions, the net  
5 effect of which does not result in an adverse functional dissimilarity between the reference and subject sequences. Typically, such a substantially equivalent sequence varies from one of those listed herein by no more than about 20%, i.e., the number of individual residue substitutions, additions, and/or deletions in a substantially equivalent  
10 sequence, as compared to the corresponding reference sequence, divided by the total number of residues in the substantially equivalent sequence is about 0.2 or less. Such a sequence is said to have 80% sequence identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified by applying the foregoing  
15 algorithm.

In one embodiment, a substantially equivalent, e.g., mutant, sequence of the invention varies from a listed sequence by no more than 10%, i.e., the number of individual residue substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the  
20 corresponding reference sequence, divided by the total number of residues in the substantially equivalent sequence is about 0.1 or less. Such a sequence is said to have 90% sequence identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified by applying the foregoing algorithm.

25 In an alternate embodiment a substantially equivalent sequence of the invention varies from a listed sequence by no more than by no more than 5%, i.e., the number of individual residue substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number  
30 of residues in the substantially equivalent sequence is about 0.05 or less. Such a sequence is said to have 95% sequence identity to the listed sequence. Such a substantially



equivalent sequence can be routinely identified by applying the foregoing algorithm.

In yet another alternate embodiment, a substantially equivalent sequence of the invention varies from a listed  
5 sequences by no more than 2%, i.e., the number of individual residue substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of residues in the substantially equivalent sequence is about 0.02 or less. Such a sequence is said to have 98% sequence  
10 identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified by applying the foregoing algorithm.

Substantially equivalent, e.g., mutant, amino acid sequences according to the invention generally have at least 95% sequence identity with a listed amino acid sequence,  
15 whereas substantially equivalent nucleotide sequence of the invention can have lower percent sequence identities, taking into account, for example, the redundancy or degeneracy of the genetic code. For the purposes of the present invention, sequences having substantially equivalent biological activity and substantially equivalent expression characteristics are  
20 considered substantially equivalent. In a preferred embodiment, for the purposes of determining equivalence, truncation of the mature sequence (e.g., via a mutation which creates a spurious stop codon) are disregarded.

Nucleic acid sequences encoding such substantially equivalent sequences, e.g., sequences of the recited percent  
25 identities, can also routinely be isolated and identified via standard hybridization procedures well known to those of skill in the art.

Where desired, an expression vector may be designed to contain a "signal or leader sequence" which will direct the polypeptide through the membrane of a cell. Such a sequence  
30 may be naturally present on the polypeptides of the present invention or provided from heterologous protein sources by recombinant DNA techniques.

A polypeptide "fragment," "portion," or "segment" is a stretch of amino acid residues of at least about 5 amino acids, often at least about 7 amino acids or about at least about 9 to 13 amino acids, and, in various embodiments, at least about 17, 25, 50, 75, 100, 150, 200, 300, 400 or more amino acids. To be "active," any polypeptide must have sufficient length to display biologic and/or immunologic activity.

Recombinant variants encoding these same or similar polypeptides may be synthesized or selected by making use of the "redundancy" in the genetic code. Various codon substitutions, such as the silent changes which produce various restriction sites, may be introduced to optimize cloning into a plasmid or viral vector or expression in a particular prokaryotic or eukaryotic system. Mutations in the polynucleotide sequence may be reflected in the polypeptide or domains of other peptides added to the polypeptide to modify the properties of any part of the polypeptide, to change characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover rate. Such variant nucleic acids and polypeptides are to be considered part of the present invention.

20       The term "activated" cells as used herein refers to those cells that are engaged in extracellular or intracellular membrane trafficking, including the export of neurosecretory or enzymatic molecules as part of a normal or disease process.

The term "purified" as used herein denotes that the indicated nucleic acid or polypeptide is present in the substantial absence of other biological macromolecules, e.g., polynucleotides, proteins, and the like. In one embodiment, the polynucleotide or polypeptide is purified such that it constitutes at least 95% by weight, more preferably at least 99.8% by weight, of the indicated biological macromolecules present (but water, buffers, and other small molecules, especially molecules having a molecular weight of less than 1000 daltons, can be present).

[illegible]

The term "secreted" protein refers to a protein that is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence when it is expressed in a suitable host cell. "Secreted" proteins include without limitation proteins secreted wholly (e.g., soluble proteins) or partially (e.g., receptors, including seven-transmembrane receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins which are transported across the membrane of the endoplasmic reticulum. Each of the above terms is meant to encompass all that is described for each, unless the context dictates otherwise.

NY2 - 924563.1

000541 034300

Nucleotide and amino acid sequences of the invention are reported below. Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of  
5 the protein may be in linear form or they may be cyclized using known methods, for example, as described in H. U. Saragovi, et al., Bio/Technology 10, 773-778 (1992) and in R. S. McDowell, et al., J. Amer. Chem. Soc. 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as  
10 immunoglobulins for many purposes, including increasing the valency of protein binding sites. For example, fragments of the protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin. For a bivalent form of the protein, such a fusion could be to the Fc portion of an IgG molecule. Other immunoglobulin isotypes may also be used to  
15 generate such fusions. For example, a protein--IgM fusion would generate a decavalent form of the protein of the invention.

The present invention also provides both full-length and mature forms of the disclosed proteins. The full-length forms of the polypeptides of the invention are identified in the  
20 figures and the sequence listing by translation of the nucleotide sequence of each nucleic acid molecule. Mature forms of the polypeptides of the invention can routinely be obtained by expression of the disclosed nucleotides encoding the full-length polypeptides in a suitable mammalian cell or other host cell. The sequence of the mature forms of  
25 the polypeptides can also routinely be determined from the amino acid sequence of the full-length polypeptides.

The present invention also provides genes corresponding to cDNA sequences disclosed herein. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include  
30 the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials.



polypeptide comprising the amino acid sequence of SEQ ID NO:2 (also referred to herein as "CD39L2"); or a polynucleotide encoding a polypeptide comprising amino acid residues 72-93, 147-162, 191-211 OR 217-238 of SEQ ID NO:2.

- 5 In selected embodiments, such isolated polynucleotides of the invention represents a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1; or a polynucleotide comprising nucleotides 232-1599, 445-510, 670-717, 802-864 or 880-945 of the nucleotide sequence of SEQ ID NO:1.

- The polynucleotides of the present invention also  
10 include, but are not limited to, a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NO:1 under highly stringent hybridization conditions; a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NO:1 under moderately stringent hybridization conditions; a polynucleotide which is an  
15 allelic variant of any polynucleotide recited above; a polynucleotide which encodes a species homologue of any of the proteins recited above; or a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptide of SEQ ID NO:2. Such polynucleotides hybridize under the above conditions to the complement of SEQ  
20 ID NO:1 or to a fragment of SEQ ID NO:1, wherein the fragment is greater than at least about 10 bp, and, in alternate embodiments, is about 20 to about 50 bp, or is greater than about 100 bp, 200 bp, 300 bp, 400 bp, 500 bp, 600 bp, 700 bp, or 800 bp.

- The isolated polynucleotides of the invention still  
25 further include, but are not limited to, a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:4 (also referred to herein as "CD39L3"); or a polynucleotide encoding a polypeptide comprising amino acid residues 55-76, 132-150, 179-199 or 213-234 of SEQ ID NO:4.

- In selected embodiments, such isolated polynucleotides  
30 of the invention represents a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3; or a polynucleotide comprising nucleotides 83-1669, 245-310, 476-532, 611-679 or 719-784 of the nucleotide sequence of SEQ ID NO:3.



nucleotide sequence of SEQ ID NO:5 under moderately stringent hybridization conditions; a polynucleotide which is an allelic variant of any polynucleotide recited above; a polynucleotide which encodes a species homologue of any of  
5 the proteins recited above; or a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptide of SEQ ID NO:6. Such polynucleotides hybridize under the above conditions to the complement of SEQ ID NO:5 or to a fragment of SEQ ID NO:5, wherein the fragment is greater than at least about 10 bp, and, in alternate  
10 embodiments, is about 20 to about 50 bp, or is greater than about 100 bp, 200 bp, 300 bp, 400 bp, 500 bp, 600 bp, 700 bp or 800 bp.

The isolated polynucleotides of the invention further include, but are not limited to, a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:8  
15 (also referred to herein as "mCD39L4" or "mNTPase"); or a polynucleotide encoding amino acid residues 46-67, 122-140, 166-187 or 194-213 of SEQ ID NO:8.

In selected embodiments, such isolated polynucleotides of the invention represents a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7; or a polynucleotide  
20 comprising nucleotides 205-1599, 340-395, 568-624, 700-765 or 784-843 of the nucleotide sequence of SEQ ID NO:7.

The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NO:7 under highly stringent hybridization conditions;  
25 a polynucleotide that hybridizes to the complement of the nucleotide sequence of SEQ ID NO:7 under moderately stringent hybridization conditions; a polynucleotide which is an allelic variant of any polynucleotide recited above; a polynucleotide which encodes a species homologue of any of the proteins recited above; or a polynucleotide that encodes  
30 a polypeptide comprising a specific domain or truncation of the polypeptide of SEQ ID NO:8. Such polynucleotides hybridize under the above conditions to the complement of SEQ ID NO:7 or to a fragment of SEQ ID NO:7, wherein the fragment



is greater than at least about 10 bp, and, in alternate embodiments, is about 20 to about 50 bp, or is greater than about 100 bp, 200 bp, 300 bp, 400 bp, 500 bp, 600 bp, 700 bp or 800 bp.

- 5       The polynucleotides of the invention additionally include the complement of any of the polynucleotides recited above.

The polynucleotides of the invention also provide polynucleotides that are substantially equivalent to the polynucleotides recited above. Typically, such a

- 10 substantially equivalent sequence varies from one of those listed herein by no more than about 20%, i.e., the number of individual nucleotide substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of nucleotides in the substantially equivalent  
15 sequence is about 0.2 or less. Such a sequence is said to have 80% sequence identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified by applying the foregoing algorithm.

- In one embodiment, a substantially equivalent polynucleotide sequence of the invention varies from a listed  
20 sequence by no more than 10%, i.e., the number of individual nucleotide substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of nucleotides in the substantially equivalent sequence is about 0.1 or less. Such a sequence is said to have 90%  
25 sequence identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified by applying the foregoing algorithm.

- In an alternate embodiment a substantially equivalent sequence of the invention varies from a listed sequence by no more than by no more than 5%, i.e., the number of individual  
30 nucleotide substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of nucleotides in the substantially equivalent sequence is

about 0.05 or less. Such a sequence is said to have 95% sequence identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified by applying the foregoing algorithm.

- 5 In yet another alternate embodiment, a substantially equivalent sequence of the invention varies from a listed sequences by no more than 2%, i.e., the number of individual nucleotide substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number
- 10 of nucleotides in the substantially equivalent sequence is about 0.02 or less. Such a sequence is said to have 98% sequence identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified.

- A polynucleotide according to the invention can be
- 15 joined to any of a variety of other nucleotide sequences by well-established recombinant DNA techniques (see Sambrook J et al. (1989) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, NY). Useful nucleotide sequences for joining to polypeptides include an assortment of vectors, e.g., plasmids, cosmids, lambda phage derivatives, phagemids,
- 20 and the like, that are well known in the art. Accordingly, the invention also provides a vector including a polynucleotide of the invention and a host cell containing the polynucleotide. In general, the vector contains an origin of replication functional in at least one organism, convenient restriction endonuclease sites, and a selectable
- 25 marker for the host cell. Vectors according to the invention include expression vectors, replication vectors, probe generation vectors, and sequencing vectors. A host cell according to the invention can be a prokaryotic or eukaryotic cell and can be a unicellular organism or part of a multicellular organism.

- 30 The sequences falling within the scope of the present invention are not limited to the specific sequences herein described, but also include allelic variations thereof. Allelic variations can be routinely determined by comparing





Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), a-factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM 1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced or derepressed by appropriate means

Page 20 of 20

(e.g., temperature shift or chemical induction) and cells are cultured for an additional period. Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for  
5 further purification.

The nucleic acid sequences of the invention are further directed to sequences which encode variants of the described nucleic acids. These amino acid sequence variants may be prepared by methods known in the art by introducing appropriate nucleotide changes into a native or variant  
10 polynucleotide. There are two variables in the construction of amino acid sequence variants: the location of the mutation and the nature of the mutation. The amino acid sequence variants of the nucleic acids are preferably constructed by mutating the polynucleotide to give an amino acid sequence that does not occur in nature. These amino  
15 acid alterations can be made at sites that differ in the nucleic acids from different species (variable positions) or in highly conserved regions (constant regions). Sites at such locations will typically be modified in series, e.g., by substituting first with conservative choices (e.g., hydrophobic amino acid to a different hydrophobic amino acid)  
20 and then with more distant choices (e.g., hydrophobic amino acid to a charged amino acid), and then deletions or insertions may be made at the target site. Amino acid sequence deletions generally range from about 1 to 30 residues, preferably about 1 to 10 residues, and are typically contiguous. Amino acid insertions include amino-  
25 and/or carboxyl-terminal fusions ranging in length from one to one hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Intrasequence insertions may range generally from about 1 to 10 amino residues, preferably from 1 to 5 residues. Examples  
30 of terminal insertions include the heterologous signal sequences necessary for secretion or for intracellular targeting in different host cells.

In a preferred method, polynucleotides encoding the novel nucleic acids are changed via site-directed mutagenesis. This method uses oligonucleotide sequences that encode the polynucleotide sequence of the desired amino acid variant, as well as a sufficient adjacent nucleotide on both sides of the changed amino acid to form a stable duplex on either side of the site of being changed. In general, the techniques of site-directed mutagenesis are well known to those of skill in the art and this technique is exemplified by publications such as, Edelman et al., DNA 2:183 (1983). A versatile and efficient method for producing site-specific changes in a polynucleotide sequence was published by Zoller and Smith, Nucleic Acids Res. 10:6487-6500 (1982). PCR may also be used to create amino acid sequence variants of the novel nucleic acids. When small amounts of template DNA are used as starting material, primer(s) that differs slightly in sequence from the corresponding region in the template DNA can generate the desired amino acid variant. PCR amplification results in a population of product DNA fragments that differ from the polynucleotide template encoding the polypeptide at the position specified by the primer. The product DNA fragments replace the corresponding region in the plasmid and this gives the desired amino acid variant.

A further technique for generating amino acid variants is the cassette mutagenesis technique described in Wells et al., Gene 34:315 (1985); and other mutagenesis techniques well known in the art, such as, for example, the techniques in Sambrook et al., supra, and Current Protocols in Molecular Biology, Ausubel et al. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be used in the practice of the invention for the cloning and expression of these novel nucleic acids. Such DNA sequences include those which are capable of hybridizing to the appropriate novel nucleic acid sequence under stringent conditions.

Finally, it is to be understood that the nucleic acid molecules of the invention further include any nucleic acid molecule that encodes the polypeptides of the invention, as described in Section 5.4, below.

5

### 5.3. Hosts

The present invention further provides host cells genetically engineered to contain the polynucleotides of the invention. For example, such host cells may contain nucleic acids of the invention introduced into the host cell using  
10 known transformation, transfection or infection methods. The present invention still further provides host cells genetically engineered to express the polynucleotides of the invention, wherein such polynucleotides are in operative association with a regulatory sequence heterologous to the host cell which drives expression of the polynucleotides in  
15 the cell.

The host cell can be a higher eukaryotic host cell, such as a mammalian cell, a lower eukaryotic host cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the recombinant construct into the host cell can be effected by calcium  
20 phosphate transfection, DEAE, dextran mediated transfection, or electroporation (Davis, L. et al., *Basic Methods in Molecular Biology* (1986)). The host cells containing one of polynucleotides of the invention, can be used in conventional manners to produce the gene product encoded by the isolated fragment (in the case of an ORF) or can be used to produce a  
25 heterologous protein under the control of the EMF.

Any host/vector system can be used to express one or more of the ORFs of the present invention. These include, but are not limited to, eukaryotic hosts such as HeLa cells, Cv-1 cell, COS cells, and Sf9 cells, as well as prokaryotic host such as *E. coli* and *B. subtilis*. The most preferred  
30 cells are those which do not normally express the particular polypeptide or protein or which expresses the polypeptide or protein at low natural level. Mature proteins can be



FOIA b 7 - D

expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., in *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, New York (1989), the disclosure of which is hereby incorporated by reference.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, *Cell* 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements. Recombinant polypeptides and proteins produced in bacterial culture are usually isolated by initial extraction from cell pellets, followed by one or more salting-out, aqueous ion exchange or size exclusion chromatography steps. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps. Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

A number of types of cells may act as suitable host cells for expression of the protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431  
5 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

Alternatively, it may be possible to produce the protein  
10 in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus*  
15 *subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be  
20 accomplished using known chemical or enzymatic methods.

In another embodiment of the present invention, cells and tissues may be engineered to express an endogenous gene comprising the polynucleotides of the invention under the control of inducible regulatory elements, in which case the regulatory sequences of the endogenous gene may be replaced  
25 by homologous recombination. As described herein, gene targeting can be used to replace a gene's existing regulatory region with a regulatory sequence isolated from a different gene or a novel regulatory sequence synthesized by genetic engineering methods. Such regulatory sequences may be comprised of promoters, enhancers, scaffold-attachment  
30 regions, negative regulatory elements, transcriptional initiation sites, regulatory protein binding sites or combinations of said sequences. Alternatively, sequences which affect the structure or stability of the RNA or protein

Patent 5,272,071

produced may be replaced, removed, added, or otherwise modified by targeting, including polyadenylation signals. mRNA stability elements, splice sites, leader sequences for enhancing or modifying transport or secretion properties of the protein, or other sequences which alter or improve the function or stability of protein or RNA molecules.

The targeting event may be a simple insertion of the regulatory sequence, placing the gene under the control of the new regulatory sequence, e.g., inserting a new promoter or enhancer or both upstream of a gene. Alternatively, the targeting event may be a simple deletion of a regulatory element, such as the deletion of a tissue-specific negative regulatory element. Alternatively, the targeting event may replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally occurring elements. Here, the naturally occurring sequences are deleted and new sequences are added. In all cases, the identification of the targeting event may be facilitated by the use of one or more selectable marker genes that are contiguous with the targeting DNA, allowing for the selection of cells in which the exogenous DNA has integrated into the host cell genome. The identification of the targeting event may also be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively selectable marker is linked to the exogenous DNA, but configured such that the negatively selectable marker flanks the targeting sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result in the stable integration of the negatively selectable marker. Markers useful for this purpose include the Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial xanthine-guanine phosphoribosyl-transferase (gpt) gene.

The gene targeting or gene activation techniques which can be used in accordance with this aspect of the invention are more particularly described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.;

International Application No. PCT/US92/09627 (WO93/09222) by Selden et al.; and International Application No. PCT/US90/06436 (WO91/06667) by Skoultchi et al., each of which is incorporated by reference herein in its entirety.

5

#### 5.4. Polypeptides of the Invention

The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising the amino acid sequence of SEQ ID NO:2; or a polypeptide comprising amino acid residues 72-93, 147-162, 191-211 OR 217-238 of SEQ ID

10 NO:2.

The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising the amino acid sequence of SEQ ID NO:4; or a polypeptide comprising amino acid residues 55-76, 132-150, 179-199 or 213-234 of SEQ ID NO:4.

15 The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising the amino acid sequence of SEQ ID NO:6; or a polypeptide comprising amino acid residues 47-68, 123-138, 167-187 or 193-214 of SEQ ID NO:6.

20 The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising the amino acid sequence of SEQ ID NO:8; or a polypeptide comprising amino acid residues 46-67, 122-140, 166-187 or 194-213 of SEQ ID NO:8.

25 The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising the amino acid sequence of SEQ ID NO:9; or a polypeptide comprising amino acid residues 77-98, 153-167, 197-217 or 223-242 of SEQ ID NO:9.

The isolated polypeptides of the invention further include polypeptides that are substantially equivalent to the polypeptides of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:9 or to specific domains thereof.

30 Typically, such a substantially equivalent sequence varies from one of those listed herein by no more than about 20%, i.e., the number of individual amino acid residue

substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of amino acid residues in the substantially equivalent sequence is about 0.2 or less. Such a sequence is said to have 80% sequence identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified by applying the foregoing algorithm.

In one embodiment, a substantially equivalent polypeptide sequence of the invention varies from a listed sequence by no more than 10%, i.e., the number of individual amino acid substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of amino acid residues in the substantially equivalent sequence is about 0.1 or less. Such a sequence is said to have 90% sequence identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified by applying the foregoing algorithm.

In an alternate embodiment a substantially equivalent sequence of the invention varies from a listed sequence by no more than by no more than 5%, i.e., the number of individual amino acid substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of amino acid residues in the substantially equivalent sequence is about 0.05 or less. Such a sequence is said to have 95% sequence identity to the listed sequence. Such a substantially equivalent sequence can be routinely identified by applying the foregoing algorithm.

In yet another alternate embodiment, a substantially equivalent sequence of the invention varies from a listed sequences by no more than 2%, i.e., the number of individual amino acid residue substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of amino acid residues in the substantially equivalent sequence is about 0.02 or less. Such a sequence is said to









FILED IN 25060

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and may thus be useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are believed to be encompassed by the present invention.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, Calif., U.S.A. (the MaxBat.RTM. kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (i.e., from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl.RTM. or Cibacrom blue 3GA Sepharose.RTM.; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression

and purification of such fusion proteins are commercially available from New England BioLab (Beverly, Mass.), Pharmacia (Piscataway, N.J.) and In Vitrogen, respectively. The protein can also be tagged with an epitope and subsequently purified  
 5 by using a specific antibody directed to such epitope. One such epitope ("Flag") is commercially available from Kodak (New Haven, Conn.).

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or  
 10 other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the  
 15 present invention as an "isolated protein."

It is to be understood that polypeptides consisting of the following amino acid sequence are not considered part of the present invention: the amino acid sequence of, or, where appropriate, reported to be encoded by the nucleotide sequence of Genbank™ accession No :S73813, gil1754710,  
 20 U91510, U91511, AA116990, AA120757, HO8436, AA378537, AA336644, AA338117, AA337885, N72742, AA256016, AA611283, AA647051, AA638277, AA271520, W46136, AA391695, AA390461, AA201196, AA246996 or AA567512.

#### 5.5. Uses And Biological Activity

25 The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration  
 30 or use of polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA).

#### 5.5.1. Research Uses and Utilities

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant

5 protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to

10 identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process

15 of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide

20 encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris et al., Cell 75:791-803 (1993)) to identify polynucleotides encoding the other protein with which binding

25 occurs or to identify inhibitors of the binding interaction.

The proteins provided by the present invention can similarly be used in assay to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in

30 assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a

particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a  
5 receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

10 Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A  
15 Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

#### 20 5.5.2. Nutritional Uses

Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the  
25 protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be added to the medium in or on which the  
30 microorganism is cultured.

5.5.3. Cytokine and Cell Proliferation/  
Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+(preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli, et al., I. Immunol. 149:3778-3783, 1992; Bowman et al., I. Immunol. 152:1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A. M. and Shevach, E. M. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human interleukin .gamma., Schreiber, R. D. In Current Protocols in Immunology. J. E.

e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without

- 5 limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L. S. and Lipsky, P. E. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988;
- 10 Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6--Nordan, R. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11--
- 15 Bennett, F., Giannotti, J., Clark, S. C. and Turner, K. J. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9--Ciarletta, A., Giannotti, J., Clark, S. C. and Turner, K. J. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.13.1, John
- 20 Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: Current Protocols in

- 25 Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA
- 30 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

5.5.4. Immune Stimulating or Suppressing Activity

A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are  
5 described herein. A polynucleotide of the invention can encode a polypeptide exhibiting such activities. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and  
10 proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by vital (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of  
15 the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, a protein of the present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the  
20 treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune  
25 thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune  
30 suppression is desired (including, for example, organ transplantation), may also be treatable using a protein of the present invention.

Using the proteins of the invention it may also be possible to immune responses, in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent.

10 Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence

15 of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as, for example, B7)), e.g., preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ

20 transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The

25 administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (e.g., B7-1, B7-3) or blocking

30 antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter





blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, 5 systemic lupus erythmatosis in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B 10 lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in 15 cases of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-vital immune responses may be enhanced in an infected patient by removing T cells from the 20 patient, costimulating the T cells in vitro with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the in vitro activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to 25 isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and 30 thereby activate, T cells in vivo.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary

- costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient mounts of
- 5 MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I  $\alpha$  .alpha. chain protein and .beta..sub.2 microglobulin protein or an MHC class II .alpha. chain protein and an MHC class II .beta. chain protein to thereby express MHC class I or MHC
  - 10 class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an
  - 15 MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to
  - 20 overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

- Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M.
- 25 Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J.
  - 30 Immunol. 135:1564-1572, 1985; Takai et al., I. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982;

- Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., Cellular Immunology 133:327-341, 1991;
- 5 Brown et al., J. Immunol. 153:3079-3092, 1994.

- Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: In vitro antibody production, Mond, J. J. and Brunswick, M. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.
- 10

- Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans);
- 15
- 20 Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

- Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995; Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994;
- 25
- 30 Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.



the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually  
5 treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either in-vivo or ex-vivo (i.e., in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or  
10 heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

15 Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. Cellular Biology 15:141-151, 1995; Keller et al., Molecular and Cellular Biology 13:473-486, 1993; McClanahan et al., Blood  
20 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lymphohematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M. G. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds.  
25 Vol pp. 265-268, Wiley-Liss, Inc., New York, N.Y. 1994; Hirayama et al., Proc. Natl. Acad. Sci. USA 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I. K. and Briddell, R. A. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, N.Y. 1994;  
30 Neben et al., Experimental Hematology 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R. E. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, N.Y. 1994; Long

term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, N.Y. 1994; Long term culture  
5 initiating cell assay, Sutherland, H. J. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, N.Y. 1994.

#### 5.5.6. Tissue Growth Activity

A protein of the present invention also may have utility  
10 in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns, incisions and ulcers.

A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is  
15 not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an  
20 osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to attract  
25 bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase  
30 activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein of the present invention

is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or  
5 ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing  
10 defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons  
15 or ligaments. The compositions of the present invention may provide environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors ex vivo for return in vivo to effect tissue  
20 repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein of the present invention may also be useful  
25 for proliferation of neural cells and for regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the  
30 treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's



100-20-20000

disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. 5 Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including 10 without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, 15 pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention 20 may also exhibit angiogenic activity.

A protein of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A protein of the present invention may also be useful 25 for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for tissue generation activity include, without 30 limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve,

neuronal); International Patent Publication No. WO91/07491 (skin, endothelium).

Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, H. I. and Rovee, D. T., eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

#### 5.5.7. Activin/Inhibin Activity

A protein of the present invention may also exhibit  
10 activin- or inhibin-related activities. A polynucleotide of the invention may encode a polypeptide exhibiting such characteristics. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins and are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH).  
15 Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin  $\alpha$ -family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these  
20 mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- $\beta$  group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, U.S. Pat. No. 4,798,885. A protein of the  
25 invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

30 Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., Endocrinology 91:562-572, 1972; Ling et al., Nature 321:779-782, 1986; Vale et al., Nature 321:776-779, 1986; Mason et al., Nature

318:659-663, 1985; Forage et al., Proc. Natl. Acad. Sci. USA  
83:3091-3095, 1986.

#### 5.5.8. Chemotactic/Chemokinetic Activity

5 A protein of the present invention may have chemotactic  
or chemokinetic activity (e.g., act as a chemokine) for  
mammalian cells, including, for example, monocytes,  
fibroblasts, neutrophils, T-cells, mast cells, eosinophils,  
epithelial and/or endothelial cells. A polynucleotide of the  
invention can encode a polypeptide exhibiting such  
10 attributes. Chemotactic and chemokinetic proteins can be  
used to mobilize or attract a desired cell population to a  
desired site of action. Chemotactic or chemokinetic proteins  
provide particular advantages in treatment of wounds and  
other trauma to tissues, as well as in treatment of localized  
infections. For example, attraction of lymphocytes, monocytes  
15 or neutrophils to tumors or sites of infection may result in  
improved immune responses against the tumor or infecting  
agent.

A protein or peptide has chemotactic activity for a  
particular cell population if it can stimulate, directly or  
indirectly, the directed orientation or movement of such cell  
20 population. Preferably, the protein or peptide has the  
ability to directly stimulate directed movement of cells.  
Whether a particular protein has chemotactic activity for a  
population of cells can be readily determined by employing  
such protein or peptide in any known assay for cell  
chemotaxis.

25 The activity of a protein of the invention may, among  
other means, be measured by the following methods:

Assays for chemotactic activity (which will identify  
proteins that induce or prevent chemotaxis) consist of assays  
that measure the ability of a protein to induce the migration  
of cells across a membrane as well as the ability of a  
30 protein to induce the adhesion of one cell population to  
another cell population. Suitable assays for movement and  
adhesion include, without limitation, those described in:  
Current Protocols in Immunology, Ed by J. E. Coligan, A. M.

- Kruisbeek, D. H. Marguiles, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995;
- 5 Lind et al. APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25:1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153:1762-1768, 1994.

#### 5.5.9. Hemostatic and Thrombolytic Activity

- 10 A protein of the invention may also exhibit hemostatic or thrombolytic activity. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other
- 15 hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

- 20 The activity of a protein of the invention may, among other means, be measured by the following methods:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79

25 (1991); Schaub, Prostaglandins 35:467-474, 1988.

#### 5.5.10. Receptor/Ligand Activity

- A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. A polynucleotide
- 30 of the invention can encode a polypeptide exhibiting such characteristics. Examples of such receptors and ligands include, without limitation, cytokine receptors and their

ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, 5 integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the 10 present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include 15 without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; 20 Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

#### 5.5.11. Anti-Inflammatory Activity

25 Proteins of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the 30 inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities

can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation intimation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)),  
5 ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to  
10 an antigenic substance or material.

#### 5.5.12. Leukemias

Leukemias and related disorders may be treated or prevented by administration of a therapeutic that promotes or inhibits function of the polynucleotides and/or polypeptides  
15 of the invention. Such leukemias and related disorders include but are not limited to acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia, chronic leukemia, chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia (for  
20 a review of such disorders, see Fishman et al., 1985, *Medicine*, 2d Ed., J.B. Lippincott Co., Philadelphia).

#### 5.5.13. Nervous System Disorders

Nervous system disorders, involving cell types which can be tested for efficacy of intervention with compounds that  
25 modulate the activity of the polynucleotides and/or polypeptides of the invention, and which can be treated upon thus observing an indication of therapeutic utility, include but are not limited to nervous system injuries, and diseases or disorders which result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination.  
30 Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients) according to the invention include but are not limited to the following

lesions of either the central (including spinal cord, brain)  
or peripheral nervous systems:

- 5 (i) traumatic lesions, including lesions caused by  
physical injury or associated with surgery,  
for example, lesions which sever a portion of  
the nervous system, or compression injuries;
- 10 (ii) ischemic lesions, in which a lack of oxygen in  
a portion of the nervous system results in  
neuronal injury or death, including cerebral  
infarction or ischemia, or spinal cord  
infarction or ischemia;
- 15 (iii) infectious lesions, in which a portion of the  
nervous system is destroyed or injured as a  
result of infection, for example, by an  
abscess or associated with infection by human  
immunodeficiency virus, herpes zoster, or  
herpes simplex virus or with Lyme disease,  
tuberculosis, syphilis;
- 20 (iv) degenerative lesions, in which a portion of  
the nervous system is destroyed or injured as  
a result of a degenerative process including  
but not limited to degeneration associated  
with Parkinson's disease, Alzheimer's disease,  
Huntington's chorea, or amyotrophic lateral  
sclerosis;
- 25 (v) lesions associated with nutritional diseases  
or disorders, in which a portion of the  
nervous system is destroyed or injured by a  
nutritional disorder or disorder of metabolism  
including but not limited to, vitamin B12  
deficiency, folic acid deficiency, Wernicke  
disease, tobacco-alcohol amblyopia,  
Marchiafava-Bignami disease (primary  
degeneration of the corpus callosum), and  
30 alcoholic cerebellar degeneration;
- (vi) neurological lesions associated with systemic  
diseases including but not limited to diabetes  
(diabetic neuropathy, Bell's palsy), systemic

lupus erythematosus, carcinoma, or sarcoidosis;

(vii) lesions caused by toxic substances including alcohol, lead, or particular neurotoxins; and

5 (viii) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including but not limited to multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

10  
Therapeutics which are useful according to the invention for treatment of a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not  
15 by way of limitation, therapeutics which elicit any of the following effects may be useful according to the invention:

(i) increased survival time of neurons in culture;  
(ii) increased sprouting of neurons in culture or *in vivo*;

20 (iii) increased production of a neuron-associated molecule in culture or *in vivo*, e.g., choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or

(iv) decreased symptoms of neuron dysfunction *in vivo*.

25 Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may be measured by the method set forth in Arakawa et al. (1990, J. Neurosci. 10:3507-3515); increased sprouting of neurons may be detected by methods set forth in Pestronk et al. (1980, Exp. Neurol. 70:65-82) or Brown et al. (1981, Ann. Rev. Neurosci. 4:17-42); increased production of neuron-  
30 associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, etc., depending on the molecule to be measured; and motor neuron dysfunction



may be measured by assessing the physical manifestation of motor neuron disorder, e.g., weakness, motor neuron conduction velocity, or functional disability.

In a specific embodiments, motor neuron disorders  
5 that may be treated according to the invention include but are not limited to disorders such as infarction, infection, exposure to toxin, trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as disorders that selectively affect neurons such as amyotrophic  
10 lateral sclerosis, and including but not limited to progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary Motorsensory Neuropathy (Charcot-  
15 Marie-Tooth Disease).

#### 5.5.14. Other Activities

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious  
20 agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation  
25 or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, co-factors or other  
30 nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders),

depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages;  
5 hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen  
10 in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

#### **5.6. Pharmaceutical Formulations and Routes of Administration**

15 A protein of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources) may be administered to a patient in need, by itself, or in pharmaceutical compositions where it is mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a variety of disorders. Such a  
20 composition may also contain (in addition to protein and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The  
25 characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-  
30 CSF, thrombopoietin, stem cell factor, and erythropoietin. The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or compliment its activity or use in treatment. Such additional

100-434500  
SECRET

factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein of the invention, or to minimize side effects. Conversely, protein of the present invention may be included in

5 formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent. A protein of the present invention may be active in

10 multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

Techniques for formulation and administration of the compounds of the instant application may be found in

15 "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition. A therapeutically effective dose further refers to that amount of the compound sufficient to result in amelioration of symptoms, e.g., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or

20 amelioration of such conditions. When applied to an individual active ingredient, administered alone, a therapeutically effective dose refers to that ingredient alone. When applied to a combination, a therapeutically effective dose refers to combined amounts of the active ingredients that result in the therapeutic effect, whether

25 administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein of the present invention is administered to a mammal having a condition to be treated. Protein of the present invention may be administered in accordance with the method

30 of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic



5.6.2. Compositions/Formulations

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. These pharmaceutical compositions may be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen. When a therapeutically effective amount of protein of the present invention is administered orally, protein of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein of the present invention, and preferably from about 1 to 50% protein of the present invention.

When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present invention will be in the form of a pyrogen-free, parenterally

100-20-25000

- acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for
- 5 intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the
- 10 present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art. For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline
- 15 buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

- For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art.
- 20 Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained solid excipient, optionally grinding a resulting mixture, and processing the
- 25 mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth,
- 30 methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or

a salt thereof such as sodium alginate. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

- Pharmaceutical preparations which can be used orally
- 10 include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft
- 15 capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration. For buccal administration, the compositions may take the form of tablets
- 20 or lozenges formulated in conventional manner.

- For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane,
- 25 dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as
- 30 lactose or starch. The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection

may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain  
5   formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily  
10   injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran.  
15   Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

20   The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides. In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be  
25   administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly  
30   soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a cosolvent system comprising benzyl





starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols. Many of the proteinase inhibiting compounds of the invention may be provided as salts with pharmaceutically compatible counterions. Such

5 pharmaceutically acceptable base addition salts are those salts which retain the biological effectiveness and properties of the free acids and which are obtained by reaction with inorganic or organic bases such as sodium hydroxide, magnesium hydroxide, ammonia, trialkylamine, dialkylamine, monoalkylamine, dibasic amino acids, sodium

10 acetate, potassium benzoate, triethanol amine and the like.

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to

15 antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T

20 lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the

25 pharmaceutical composition of the invention. The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid

30 crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation

of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Pat. Nos. 4,235,871; 4,501,728; 4,837,028; and 4,737,323, all of which are incorporated herein by reference.

5       The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein of the present invention  
10 with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not  
15 increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01  $\mu$ g to about 100 mg (preferably about 0.1  $\mu$ g to about 10 mg, more preferably about 0.1  $\mu$ g to about 1 mg) of protein of the present invention per kg body weight. For compositions of the  
20 present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically  
25 acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein of the invention which may also optionally be included in the  
30 composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention. Preferably for bone and/or cartilage formation, the





and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced  
5 either in vivo or ex vivo into cells for expression in a mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA). Cells may also be cultured ex vivo in the presence of  
10 proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced in vivo for therapeutic purposes.

#### 5.6.3. Effective Dosage

15 Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the  
20 subject being treated. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For example, a dose can be  
25 formulated in animal models to achieve a circulating concentration range that includes the IC<sub>50</sub> as determined in cell culture (i.e., the concentration of the test compound which achieves a half-maximal inhibition of the C-proteinase activity). Such information can be used to more accurately determine useful doses in humans.

30 A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms or a prolongation of survival in a patient. Toxicity and

therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD<sub>50</sub> and ED<sub>50</sub>. Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED<sub>50</sub> with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. See, e.g., Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1. Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the C-proteinase inhibiting effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data; for example, the concentration necessary to achieve 50-90% inhibition of the C-proteinase using the assays described herein. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

5

#### 5.6.4. Packaging

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labelled for treatment of an indicated condition.

15

#### 5.7. Antibodies

Another aspect of the invention is an antibody that specifically binds the polypeptide of the invention. Such antibodies can be either monoclonal or polyclonal antibodies, as well fragments thereof and humanized forms or fully human forms, such as those produced in transgenic animals. The invention further provides a hybridoma that produces an antibody according to the invention. Antibodies of the invention are useful for detection and/or purification of the polypeptides of the invention.

Protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the protein. Such antibodies may be obtained using either the entire protein or fragments thereof as an immunogen. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for synthesizing such peptides are known in the art, for example, as in R. P. Merrifield, J. Amer. Chem. Soc. 85, 2149-2154 (1963); J. L. Krstenansky, et al., FEBS



SECRET

Lett. 211, 10 (1987). Monoclonal antibodies binding to the protein of the invention may be useful diagnostic agents for the immunodetection of the protein. Neutralizing monoclonal antibodies binding to the protein may also be useful

- 5 therapeutics for both conditions associated with the protein and also in the treatment of some forms of cancer where abnormal expression of the protein is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal antibodies against the protein may be useful in detecting and preventing the metastatic spread of the cancerous cells,
- 10 which may be mediated by the protein. In general, techniques for preparing polyclonal and monoclonal antibodies as well as hybridomas capable of producing the desired antibody are well known in the art (Campbell, A.M., *Monoclonal Antibodies Technology: Laboratory Techniques in Biochemistry and*
- 15 *Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1984); St. Groth et al., *J. Immunol.* 35:1-21 (1990); Kohler and Milstein, *Nature* 256:495-497 (1975)), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., *Immunology Today* 4:72 (1983); Cole et al., in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc.
- 20 (1985), pp. 77-96).

- Any animal (mouse, rabbit, etc.) which is known to produce antibodies can be immunized with a peptide or polypeptide of the invention. Methods for immunization are well known in the art. Such methods include subcutaneous or intraperitoneal injection of the polypeptide. One skilled in
- 25 the art will recognize that the amount of the protein encoded by the ORF of the present invention used for immunization will vary based on the animal which is immunized, the antigenicity of the peptide and the site of injection. The protein that is used as an immunogen may be modified or administered in an adjuvant in order to increase the
- 30 protein's antigenicity. Methods of increasing the antigenicity of a protein are well known in the art and include, but are not limited to, coupling the antigen with a







fermentation reactions and in the production of commercially useful metabolites.

As used herein, "a computer-based system" refers to the hardware means, software means, and data storage means used  
5 to analyze the nucleotide sequence information of the present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means, and data storage means. A skilled artisan can readily appreciate that any one of the currently available computer-based systems are  
10 suitable for use in the present invention. As stated above, the computer-based systems of the present invention comprise a data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means and software means for supporting and implementing a search means. As used herein, "data storage means" refers to  
15 memory which can store nucleotide sequence information of the present invention, or a memory access means which can access manufactures having recorded thereon the nucleotide sequence information of the present invention.

As used herein, "search means" refers to one or more programs which are implemented on the computer-based system  
20 to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of a known sequence which match a particular target sequence or target motif. A variety of known algorithms are disclosed publicly and a variety of commercially available  
25 software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software includes, but is not limited to, MacPattern (EMBL), BLASTN and BLASTA (NPOLYPEPTIDEIA). A skilled artisan can readily recognize that any one of the available algorithms or implementing software packages for  
30 conducting homology searches can be adapted for use in the present computer-based systems. As used herein, a "target sequence" can be any nucleic acid or amino acid sequence of six or more nucleotides or two or more amino acids. A



transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the  
5 present invention is necessary for the design of an antisense or triple helix oligonucleotide.

#### 5.10. Diagnostic Assays and Kits

The present invention further provides methods to identify the presence or expression of one of the ORFs of the  
10 present invention, or homolog thereof, in a test sample, using a nucleic acid probe or antibodies of the present invention.

In general, methods for detecting a polynucleotide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the  
15 polynucleotide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polynucleotide of the invention is detected in the sample.

Such methods can also comprise contacting a sample under stringent hybridization conditions with nucleic acid primers that anneal to a polynucleotide of the invention under such  
20 conditions, and amplifying annealed polynucleotides, so that if a polynucleotide is amplified, a polynucleotide of the invention is detected in the sample.

In general, methods for detecting a polypeptide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the polypeptide for a  
25 period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polypeptide of the invention is detected in the sample.

In detail, such methods comprise incubating a test sample with one or more of the antibodies or one or more of nucleic acid probes of the present invention and assaying for binding  
30 of the nucleic acid probes or antibodies to components within the test sample.

Conditions for incubating a nucleic acid probe or antibody with a test sample vary. Incubation conditions

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid probe or antibody used in the assay. One skilled in the art will recognize that any one of the commonly available

5 hybridization, amplification or immunological assay formats can readily be adapted to employ the nucleic acid probes or antibodies of the present invention. Examples of such assays can be found in Chard, T., *An Introduction to*

10 *Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G.R. et al., *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985);

15 Tijssen, P., *Practice and Theory of immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985). The

20 test samples of the present invention include cells, protein or membrane extracts of cells, or biological fluids such as sputum, blood, serum, plasma, or urine. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for

25 preparing protein extracts or membrane extracts of cells are well known in the art and can be readily be adapted in order to obtain a sample which is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out

25 the assays of the present invention. Specifically, the invention provides a compartment kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the probes or antibodies of the present invention; and (b) one or more other containers comprising one or more of the following: wash reagents,

30 reagents capable of detecting presence of a bound probe or antibody.

In detail, a compartment kit includes any kit in which reagents are contained in separate containers. Such



containers include small glass containers, plastic containers or strips of plastic or paper. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the antibodies used in the assay, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound antibody or probe. Types of detection reagents include labeled nucleic acid probes, labeled secondary antibodies, or in the alternative, if the primary antibody is labeled, the enzymatic, or antibody binding reagents which are capable of reacting with the labeled antibody. One skilled in the art will readily recognize that the disclosed probes and antibodies of the present invention can be readily incorporated into one of the established kit formats which are well known in the art.

#### 5.11. Screening Assays

- Using the isolated proteins and polynucleotides of the invention, the present invention further provides methods of obtaining and identifying agents which bind to a protein encoded by the ORF from a nucleic acid with a sequence of SEQ ID NO:1, to a specific domain of the polypeptide encoded by the nucleic acid, or to a nucleic acid with a sequence of SEQ ID NO:1. In detail, said method comprises the steps of:
- (a) contacting an agent with an isolated protein encoded by an ORF of the present invention, or nucleic acid of the invention; and
  - (b) determining whether the agent binds to said protein or said nucleic acid.

In general, therefore, such methods for identifying compounds that bind to a polynucleotide of the invention can comprise contacting a compound with a polynucleotide of the invention for a time sufficient to form a

polynucleotide/compound complex, and detecting the complex, so that if a polynucleotide/compound complex is detected, a compound that binds to a polynucleotide of the invention is identified.

- 5 Likewise, in general, therefore, such methods for identifying compounds that bind to a polypeptide of the invention can comprise contacting a compound with a polypeptide of the invention for a time sufficient to form a polypeptide/compound complex, and detecting the complex, so that if a polypeptide/compound complex is detected, a
- 10 compound that binds to a polynucleotide of the invention is identified.

- Methods for identifying compounds that bind to a polypeptide of the invention can also comprise contacting a compound with a polypeptide of the invention in a cell for a time sufficient to form a polypeptide/compound complex,
- 15 wherein the complex drives expression of a receptor gene sequence in the cell, and detecting the complex by detecting reporter gene sequence expression, so that if a polypeptide/compound complex is detected, a compound that binds a polypeptide of the invention is identified.

- Compounds identified via such methods can include
- 20 compounds which modulate the activity of a polypeptide of the invention (that is, increase or decrease its activity, relative to activity observed in the absence of the compound). Alternatively, compounds identified via such methods can include compounds which modulate the expression of a polynucleotide of the invention (that is, increase or
- 25 decrease expression relative to expresssion levels observed in the absence of the compound). Compounds, such as compounds identified via the methods of the invention, can be tested using standard assays well known to those of skill in the art for their ability to modulate activity/expression.

- The agents screened in the above assay can be, but are
- 30 not limited to, peptides, carbohydrates, vitamin derivatives, or other pharmaceutical agents. The agents can be selected and screened at random or rationally selected or designed using protein modeling techniques.

For random screening, agents such as peptides, carbohydrates, pharmaceutical agents and the like are selected at random and are assayed for their ability to bind to the protein encoded by the ORF of the present invention.

5 Alternatively, agents may be rationally selected or designed. As used herein, an agent is said to be "rationally selected or designed" when the agent is chosen based on the configuration of the particular protein. For example, one skilled in the art can readily adapt currently available procedures to generate peptides, pharmaceutical agents and  
10 the like capable of binding to a specific peptide sequence in order to generate rationally designed antipeptide peptides, for example see Hurby et al., Application of Synthetic Peptides: Antisense Peptides," In *Synthetic Peptides, A User's Guide*, W.H. Freeman, NY (1992), pp. 289-307, and  
15 Kaspczak et al., *Biochemistry* 28:9230-8 (1989), or pharmaceutical agents, or the like.

In addition to the foregoing, one class of agents of the present invention, as broadly described, can be used to control gene expression through binding to one of the ORFs or EMFs of the present invention. As described above, such agents can be randomly screened or rationally designed/selected. Targeting the ORF or EMF allows a skilled artisan to design sequence specific or element specific agents, modulating the expression of either a single ORF or multiple ORFs which rely on the same EMF for expression control. One class of DNA binding agents are agents which contain base residues which hybridize or form a triple helix formation by binding to DNA or RNA. Such agents can be based on the classic phosphodiester, ribonucleic acid backbone, or can be a variety of sulfhydryl or polymeric derivatives which have base attachment capacity.

Agents suitable for use in these methods usually contain 20 to 40 bases and are designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., *Nucl. Acids Res.* 6:3073 (1979); Cooney et al., *Science* 241:456 (1988); and Dervan et al., *Science*

251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988)). Triple helix- formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the present invention is necessary for the design of an antisense or triple helix oligonucleotide and other DNA binding agents. Agents which bind to a protein encoded by one of the ORFs of the present invention can be used as a diagnostic agent, in the control of bacterial infection by modulating the activity of the protein encoded by the ORF. Agents which bind to a protein encoded by one of the ORFs of the present invention can be formulated using known techniques to generate a pharmaceutical composition.

#### 5.10. Use of Nucleic Acids as Probes

Another aspect of the subject invention is to provide for polypeptide-specific nucleic acid hybridization probes capable of hybridizing with naturally occurring nucleotide sequences. The hybridization probes of the subject invention may be derived from the nucleotide sequence of the SEQ ID NO:1. Because the corresponding gene is only expressed in a limited number of tissues, especially adult tissues, a hybridization probe derived from SEQ ID NO:1 can be used as an indicator of the presence of RNA of cell type of such a tissue in a sample.

Any suitable hybridization technique can be employed, such as, for example, in situ hybridization. PCR as described US Patent Nos 4,683,195 and 4,965,188 provides additional uses for oligonucleotides based upon the nucleotide sequences. Such probes used in PCR may be of recombinant origin, may be chemically synthesized, or a mixture of both. The probe will comprise a discrete nucleotide sequence for the detection of identical sequences or a degenerate pool of

possible sequences for identification of closely related genomic sequences.

Other means for producing specific hybridization probes for nucleic acids include the cloning of nucleic acid sequences into vectors for the production of mRNA probes. Such vectors are known in the art and are commercially available and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate radioactively labeled nucleotides. The nucleotide sequences may be used to construct hybridization probes for mapping their respective genomic sequences. The nucleotide sequence provided herein may be mapped to a chromosome or specific regions of a chromosome using well known genetic and/or chromosomal mapping techniques. These techniques include in situ hybridization, linkage analysis against known chromosomal markers, hybridization screening with libraries or flow-sorted chromosomal preparations specific to known chromosomes, and the like. The technique of fluorescent in situ hybridization of chromosome spreads has been described, among other places, in Verma et al (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York NY.

Fluorescent in situ hybridization of chromosomal preparations and other physical chromosome mapping techniques may be correlated with additional genetic map data. Examples of genetic map data can be found in the 1994 Genome Issue of Science (265:1981f). Correlation between the location of a nucleic acid on a physical chromosomal map and a specific disease (or predisposition to a specific disease) may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier or affected individuals. The nucleotide sequence may be used to produce purified polypeptides using well known methods of recombinant DNA technology. Among the many publications that teach methods for the expression of genes after they have been isolated is Goeddel (1990) Gene Expression Technology, Methods and Enzymology, Vol 185,





embodiments and variations may be made in the scope of the present invention. Accordingly, it is intended that the broader aspects of the present invention not be limited to the disclosure of the following examples.

5

6. **EXAMPLE: Identification of Novel CD-39-Like  
Nucleic Acid and Polynucleotide Molecules**

Described herein is the cloning and characterization of novel CD-39-like nucleotide-triphosphatase ("NTPase") gene and polypeptide sequences. These sequences are referred to below as "mNTPase", "mCD39L4" and "CD39L2."

15  
20  
25  
30  
First, a novel murine family member was cloned by low stringency screening of mouse cDNA libraries with a human CD39L1 cDNA clone (Chadwick, B.P. & Frischauf A.-M., 1997, Mamm. Genome 8:668-672). A 1738 bp cDNA clone was isolated from an adult mouse testis cDNA library (Stratagene Ltd., Cambridge, UK) and sequenced. DNA sequence comparisons with the human CD39L1 cDNA sequence showed moderate DNA homology of approximately 39% identity). An open reading frame (ORF) could be detected for the cDNA sequence, but indicated that the cDNA clone did not contain the initiation methionine codon and, therefore, did not extend to the 5' end. Database searches with the mouse cDNA sequence identified two mouse EST clones that extended the cDNA sequence at the 5' end (Accession Nos. AA116990 and AA120757). The EST clones were resequenced. The cloned and the resequenced nucleotide sequences were analyzed and were combined appropriately to yield the nucleotide sequence (SEQ ID NO:7) depicted in FIG. 1, and referred to herein as mCD39L4 or mNTPase. The sequence revealed an ORF from nucleotides 205 to 1599 with the ATG at nucleotide 205 having a moderate match to the initiation start site for vertebrates (AAGAAUAUGG for mNTPase versus GCCGCCAUGG; Kozak, M., 1989, J. Biol. Chem. 108:229-241). The derived amino acid sequence is also shown in FIG. 2 (SEQ ID NO:8). No apparent polyadenylation signal existed, although the cDNA clone isolated contained a poly-A tail.



Hydropathy plots with Topred-II 1.1 (Claros, M.G. & Von Hejine, G., 1994, Comput. Appl. Biosci. 10:685-686) predict a single potential transmembrane segment close to the amino terminus of the protein, suggesting a single-pass  
5 transmembrane protein with a large extracellular domain. Two potential glycosylation sites can be found at amino acid positions 41 (NVSA) and at 231 (NSTF), suggesting that mNTPase is glycosylated.

Database searches with the derived amino acid sequence identified homology with other members of the NTPase family.  
10 FIG. 2 shows an alignment of the full mNTPase (mCD39L4) protein sequence against three of the most homologous known NTPases, from garden pea, potato and *Saccharomyces cerevisiae*. The mNTPase protein shares approximately 30% amino acid identity with the three other NTPases.

The region of highest homology between all members of  
15 the NTPase family is at the amino terminus of the protein. Handa & Guidotti (Handa, M. & Guidotti, G., 1996, Biochem. Biophys. Res. Commun. 218:916-923) highlighted four regions of NTPases referred to as putative apyrase-conserved regions ("ACRs"). FIG. 3 shows an alignment of ACRs I-IV. (See  
20 Section 3, Section 5, and its subsections, above, for a delineation of the amino acid residues that make up ACRs I-IV of the CD39-like polypeptides of the invention.) ACR conservation would indicate that these regions are essential for the functioning of the protein, while changes in the regions surrounding these domains can be tolerated. The presence of all four ACRs in the mNTPase (mCD39L2) indicates  
25 that mNTPase is a new member of the NTPase family.

BLAST searches with the DNA sequence of mNTPase (mCD39L4) revealed two overlapping human EST clones with 57% DNA sequence identity to portions of mNTPase (Accession Nos. H08436 and AA378537). Upon combination and analysis of the resulting sequence, an ORF was identified that showed  
30 homology to NTPases. The putative NTPase protein sequence, referred to herein as "CD39L2," is shown in FIG. 3 alongside the other NTPase protein sequences. The identification and

characterization of the full-length CD39L2 polypeptide and nucleotide sequences is described in the Example presented in Section 7, below.

To map the murine mNTPase gene, a cosmid was isolated  
5 from a mouse cosmid library, and used for fluorescence in situ hybridization (FISH). For the FISH analysis, slides with mouse metaphase chromosomes were prepared from spleens as described in Monier et al. (Monier, K. et al., 1996, Cytogenet. Cell Genet. 72:200-204). 1 microgram of mouse cosmid containing the mNTPase gene was labeled with biotin  
10 14-ATP and a Bionick Kit (GibcoBRL). DNA was purified by passage through a Sephadex G50 column and ethanol precipitated with 50 micrograms of sheared salmon sperm DNA and tRNA. 80 ng of probe was dried down with 3 micrograms of mouse Cot-1 DNA (GibcoBRL). Hybridization was carried out as described (Ragoussis, J. et al., 1992, Genomics 14:423-430).  
15 Confirmation of chromosomal location was achieved by rehybridizing the same slide with a mouse Chromosome 12-specific Starfish Paint (Cambio).

NS  
D3/ The FISH study revealed the presence of mNTPase on mouse  
Chr. 12 at chromosome band E. To confirm the location of the mNTPase gene on mouse Chr. 12, linkage analysis was carried  
20 out upon the European Collaborative Interspecific Backcross (EUCIB). PCR primers were designed to the 3' untranslated region of the mNTPase cDNA sequence and used for PCR by use of mouse genomic DNA from the two parental mouse strains, *Mus spretus* and C57BL/6. A polymorphism was detected between the two strains by SSCP analysis and was used for the mapping.  
25 (PCR conditions: 48°C 20 sec., Primer 1: CCAGACTGTAAATCTTTTGG; Primer 2: AGGGAATGTAATAAGGGTAG; conditions: 94°C 2 minutes; 35 cycles of 94°C 20 sec. 72°C 20 sec., 72°C 1 min; product size: 320 bp).

Linkage with a LOD score of 8.14 was obtained with the genetic marker D12Mit4, flanked by D12Mit149 and D12Mit238,  
30 between 31.7 cM and 36.1 cM from the top of the mouse Chr. 12 linkage group for the MIT F<sub>2</sub> Intercross. This region of mouse Chr. 12 has previously been shown to share synteny with human

Chr. 14q (DeBry, R.W. & Seldin, M.F., 1996, Genomics 33:337-351).

5            7.    **EXAMPLE: Identification and Characterization of  
Additional Novel CD39-Like Polypeptides  
and Nucleic Acid Molecules**

Described herein is the cloning and characterization of novel CD-39-like nucleotide-triphosphatase ("NTPase") gene and polypeptide sequences.

10 These sequences are referred to below as "CD39L2," "CD39L3," "CD39L4" and "dCD39L4."

7.1.    **Materials and Methods**

INDW Identification, isolation, and sequencing of cDNA clones  
for CD39L2, CD39L3, and CD39L4. The nucleotide sequence of  
15 CD39 (Accession No. S73813), CD39L1 (Accession No. U91510),  
and mNTPase (see Section 6, above) were used in TBLASTX  
searches against entries in the expressed sequence tag (EST)  
database at EMBL/GenBank, using the Bork server  
(<http://www.bork.embl-heidelberg.de/>). cDNA clones for  
homologous IMAGE EST entries were obtained from the Human  
20 Genome Mapping Project Resource Centre (HGMP, Hinxton, UK).  
DNA was prepared with QiaTip-100 (Qiagen), and the cDNA was  
sequenced by primer walking with a fluorescence labeled dye-  
terminator cycle sequencing kit according to the  
manufacturer's instructions (PRISM Ready Dye-Deoxy Terminator  
Premix from Applied Biosystems Inc.) and electrophoresed on  
25 an ABI 373 (Perkin-Elmer). Overlapping EST clones were  
identified by searching with the nucleotide sequence against  
entries in the EST database using BLAST-N  
(<http://www.ncbi.nlm.nih.gov:80/cgi-bin/BLAST/nph-blast?Jform>).

Additional IMAGE cDNA clones were ordered from HGMP if  
30 they extended the existing nucleotide sequence further 5'.  
cDNA clones corresponding to the most 5' extreme of each gene  
were identified by hybridization of radiolabeled inserts of  
IMAGE cDNA clones to a keratinocyte stem cell cDNA library, a

human adult breast epithelial cDNA library constructed using Stratagene Lambda ZAP vector, and a Jurkat cell line cDNA library in pBluescript (Dunne, J. et al., 1995, Genomics 30:207-223).

- 5     Northern analysis of members of the CD39-like gene family. cDNA clone inserts were removed by restriction digestion and separated by gel electrophoresis. Insert DNA was gel-purified and radiolabeled (Sambrook et al., 1989, supra). Radiolabeled cDNA was prehybridized at 65°C for 2 h with 20 µg of human Cot-1 DNA (GibcoBRL) and 100 µg of total  
10 human DNA (Sigma), before hybridization to Northern blots (Clontech, human multiple tissue Northern blots, Catalog No. 7760-1 and 7759-1) according to the manufacturer's instructions.

- INS  
DS  
15     Mapping of CD39L2, CD39L3, and CD39L4. Members of the CD39-like gene family were mapped in the human genome by PCR screening of the GeneBridge-4 radiation hybrid mapping panel obtained from the HGMP Resource Centre (Hinxton, UK) (Gyapay, G. et al., 1996, Hum. Mol. Genet. 5:339-346). PCR-positive radiation hybrid clones were organized into the GeneBridge-4 HGMP-RC subset order using the HGMP radiation hybrid mapping World Wide Web (WWW) site (<http://www.hgmp.mrc.ac.uk/cgi-bin/contig/contig/rhmapper.pl>), and mapping data for each  
20 gene were obtained from the Whitehead server (<http://www.genome.wi.mit.edu/cgi-bin/rhmapper.pl>). The chromosomal location for each gene was confirmed by PCR screening of the monochromosomal hybrids obtained from the HGMP Resource Centre. PCR primers were designed for the 3'  
25 untranslated region (UTR) of each gene and titrated for a unique human-specific PCR product. PCR conditions: CD39L2, Primer 1, 5'-CTGCTTGAGTGACGTCTCTG-3'; Primer 2, 5'-CACATGAGGTTCAGCTCGTG-3'; 94°C for 2 min; 38 cycles of 94°C for 20 s, 54°C for 20 s, 72°C for 20 s; 72°C for 2 min. Product size is 362 bp). CD39L3, Primer 1: 5'-  
30 GTGAAGTGGCTGCCTTCAGG-3'; Primer 2, 5'-CCTTTGACTCGGGACTCCAG-3'; 94°C for 2 min; 38 cycles of 94°C for 20 s, 56°C for 20 s, 72°C for 2 min. Product size is 281 bp). CD39L4. Primer 1, 5'-GAACTGCTGCCTAACCACTC-3'; Primer 2, 5'-

ATTGATGGGTCTTGGGATTGC-3'; 94°C 2 for min; 38 cycles of 94°C for 20 s, 56°C for 20 s; 72°C for 20 s; 72°C for 2 min. Product size is 234 bp. PCR products were analyzed by electrophoresis through 3.5% NuSieve agarose gels (Flowgen).

5

## 7.2. Results

### Isolation and Sequence Characterization of CD39L2.

Identification of partial human CD39L2 sequence was described in the Example presented in Section 6, above. The CD39L2 insert was used to isolate additional clones from a human adult breast epithelial cDNA library (ZR75), a human T-leukemia cell line J6 cDNA library (Jurkat), and a human keratinocyte stem cell cDNA library (KER). Of 23 cDNA clones that were isolated and sequenced, all but one appeared to be alternatively spliced or unspliced. Within the 2762 bp cDNA that appeared to be neither unspliced or alternatively spliced, an ORF extending to nucleotide 1600 containing ACRs I-IV was identified. Two ATG codons with a poor match to the consensus translation initiation site were found at nucleotide positions 148 and 232 (AUGUGAAUGA at 148 and ACAAGGAUGA at 232 versus consensus GCCGCCAUGG; Kozak, M., 1989, J. Biol. Chem. 108:229-241). Based on homology to mNTPase, the ATG at nucleotide position 232 is the initiation codon. (See FIG. 9 for a depiction of the CD39L2 amino acid sequence that results from translation from the upstream, position 148, start codon; such a form of CD39L2 as well as nucleotide sequences that encode this form of the polypeptide are also intended to be included as part of the present invention.) A single polyadenylation signal of AAUAAA was identified at nucleotide position 2700, 22 nucleotides 5' of the poly(A) tail of the human CD39L2 cDNA.

The nucleotide sequence (SEQ ID NO:1) and derived amino acid sequence (SEQ ID NO:2) of human CD39L2 is depicted in FIG. 4. Hydrophobicity plots using Topred-II 1.1 (Claros, M.G. & Von Hejine, G., 1994, Comput. Appl. Biosci. 10:685-686) predicted a single transmembrane segment at the N-terminal extreme of the protein, suggesting that CD39L2 has a short putative cytoplasmic tail and a large extracellular C-

106120-24250000

terminal domain (FIG. 5). There are two potential N-glycosylation sites in the predicted extracellular domain. A cAMP and cGMP-dependent protein kinase and a protein kinase-C phosphorylation site are found directly after the initiation  
5 methionine codon (nucleotide 232).

Isolation and Sequence Characterization of CD39L3. An additional novel gene and polypeptide CD39-like sequence, referred to herein as CD39L3, was also identified. BLASTN search of the NCBI EST database with the full cDNA sequence for human CD39 (Accession No. S73813) identified three EST  
10 entries for cDNA clones from an endometrial tumor library (Accession Nos. AA336644, AA338117, and AA337885), which led to the identification of an IMAGE library EST (Accession No. N72742) that was completely sequenced and had amino acid homology to CD39 and CD39L1. The insert of N72742 was used to screen the Jurkat, ZR75, and KER cDNA libraries, and a  
15 single cDNA clone was isolated from the KER library and sequenced.

A 1669-bp ORF was identified within the cDNA insert. The nucleotide sequence (SEQ ID NO:3) and derived amino acid sequence (SEQ ID NO:4) of the cDNA insert, referred to herein as CD39L3, are shown in FIG. 6. The amino acid sequence was  
20 revealed to contain ACRs I-IV, an ATG codon at position 83, and a single polyadenylation signal at position 2758. Hydrophobicity plots as described above predict two potential transmembrane segments at the N and C-terminal extremes of the protein (FIG. 5). There are seven potential extracellular N-glycosylation sites. A cAMP- and cGMP-  
25 dependent protein kinase site and a protein kinase-C phosphorylation site are located at the C-terminal extreme of the protein.

Isolation and Sequence Characterization of CD39L4. An additional CD39-like gene and polypeptide sequence, referred to herein as CD39L4, was also identified. A TBLASTX search of the NCBI EST database with the full cDNA sequence for the mNTPase was performed. A human EST clone was sequenced, and an ORF was identified extending to nucleotide 529 of 2260 nucleotides that contained ACR I only and an ATG codon at

INS  
DT  
30

position 256. In the same reading frame, downstream of the stop codon at nucleotide 529, an ORF extending to nucleotide 1792 contained ACRs II, III, and IV. Further analysis of the nucleotide sequence revealed a putative intron with splice donor and acceptor sites that conform to the 5' gt...3' ag rule (Breathnach and Chambon, 1981, Ann Rev. Biochem. 50:349-383; splice donor CAGgtcacttatggagcctg at nucleotide position 470, acceptor ccatggacaaaatagGAC at position 710, exon sequence underlined). Further analysis of the sequence revealed that removal of the 251-bp putative intron would

10 result in a contiguous ORF containing ACRs I-IV. The hypothesis that this sequence does indeed constitute an intron was only confirmed by isolation and sequencing of three additional cDNA clones (CD39LAJ1-3) from the Jurkat library, one of which contained the 251 bp.

The nucleotide sequence (SEQ ID NO:5) and derived amino acid sequence (SEQ ID NO:6), referred to herein as CD39L4, is depicted in FIG. 7. The sequence contained a poly(A)tail, but no consensus polyadenylation sequence (Proudfoot, 1991, New Biol. 3:851-854). This is also the case for mNTPase. Hydrophobicity plots as described above predicted a single transmembrane segment at the N-terminal extreme of the

20 protein (Fig. 5). This is similar to the predicted topology of CD39L2 and different from that of CD39, CD39L1, and CD39L3. There are three potential extracellular N-glycosylation sites.

FIG. 8 depicts an alignment of each of the above-described sequences, along with other members of the NTPase

25 CD-39-like gene family. The ACR domains are indicated by arrows.

#### Expression of the Human Members of the CD39-Like Family.

Representative probes for each member of the CD39-like gene family were hybridized to human multiple tissue Northern

30 blots. Results are presented below.

CD39L2. Hybridization of the CD39L2 cDNA insert to the multiple tissue Northern blots resulted in two prominent signals of 2.6 kb (major) and 4.4 kb (minor) in all tissues

studied. This is most likely due to differential polyadenylation.

CD39L3. A PCR product covering the coding sequence of CD39L3 was used for the Northern hybridizations. A strong  
5 signal of approximately 3.0 kb could be seen in adult brain, pancreas, spleen, and prostate. Though moderate or low expression was seen in most other tissues, no signal was detected in adult liver and peripheral blood leukocytes. A weaker signal of approximately 1.8 kb was found in adult pancreas and may be the result of alternative splicing.

10 CD39L4. The CD39L4 cDNA was hybridized to the same Northern blots, and a prominent signal of approximately 4.8 kb was seen in adult liver, kidney, prostate, testis, and colon. Considerably weaker expression was seen for all other tissues examined. Several smaller bands were observed in  
15 tissues showing the strongest expression of CD39L4 and may be the result of differential polyadenylation or alternative splicing.

#### Mapping of Members of the CD39-like Family.

The CD39L2 gene was mapped with a lod score of >19 to human chromosome 20 by PCR typing of the GeneBridge 4  
20 Radiation Hybrid Mapping Panel (Gyapay et al., 1996, Human Mol. Genet. 5:339-346). CD39L2 mapped 9.76 cR from D20S493 (typing data: 12012 02101 22000 00111 00110 01210 00110 01101 10121 00100 00120 11211 00011 11012 01001 01102 00000 00000 001). Using the closest flanking markers (D20S184 and D20S99) also represented on the consensus map, this placed  
25 CD39L2 at chromosome band 20q11.2. The location of CD39L2 on human chromosome 20 was confirmed by PCR analysis of monochromosomal mapping panels (Kelsell et al., 1995, Ann. Human Genet. 59:233-241). On the basis of synteny to human chromosome 20q11.2, the mouse homolog of CD39L2 was expected to map to mouse chromosome 2 (DeBry and Seldin, 1996,  
30 Genomics 33:337-351).

The CD39L3 gene was mapped as described above to human chromosome 3, 5.76 cR from D3S3390 (data: 12002 02010 22000 00011 20000 00110 01001 00000 02022 11000 10001 00200 21100



00212 01010 10002 00000 00011 001). Using the closest flanking markers as described above (D3S1561 and D3S3564), this placed CD39L3 at chromosome band 3p21.3. The location of CD39L3 on chromosome 3 was confirmed by PCR as for CD39L2.

5 On the basis of synteny, the mouse homologue of Cd39L3 was expected to map to mouse chromosome 9 (DeBry and Seldin, 1996, Genomics 33:337-351).

The CD39L4 gene was mapped as described above to human chromosome 14, 1.92 cR from D14S71 (data: 02102 02102 22000 01010 11021 01000 01010 10110 02121 21000 00010 00211 01001 10102 02012 00002 12111 01100 002). This placed CD39L4 at chromosome band 14q24. The chromosomal location of CD39L4 was confirmed as described above.

Identification of a Drosophila Gene with high Homology to CD39L2 and CD39L4. A D. melanogaster CD39-like gene was

15 also identified. A TBLASTX search of the EST database using the human CD39L2 cDNA sequence, five Drosophila EST entries were identified (Accession No. AA391695, AA390461, AA201196, AA246996, and AA567512). A consensus sequence was generated and used for a BLASTN search against EMBL/GenBank entries. A single D. melanogaster genomic entry (Accession No. AC002032)

20 was identified showing 100% sequence identity to three regions of the EST consensus sequence. Alignment of the EST consensus against the genomic sequence identified three exons that conform to the 5' gt...3' ag rule (Breathnach & Chambon, 1981, Ann. Rev. Biochem. 50:349-383). Exon 4 was identified on the basis of reading frame homology to the CD39L2 and

25 CD39L4 proteins. An ATG codon was identified in exon 1, a stop codon in exon 4.

The predicted amino acid sequence of the D. melanogaster CD39-like gene, referred to herein as dCD39L4, containing the ACRs-I-IV was shown in FIG. 9, aligned against the gene family members with the highest homology. Three N-

30 glycosylation consensus sites were found in the putative extracellular domain, and two potential cAMP- and c-GMP-dependent protein kinase phosphorylation sites were found in the putative N-terminal cytoplasmic domain. Hydrophobicity

plots as described above predicted a single transmembrane segment at the N-terminal extreme of the dCD39L4 protein. The topology of dCD39L4 is therefore most similar to the predicted topology of the CD39L2 and Cd39L4 proteins.

5

The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the invention, and compositions and methods which are functionally equivalent are within the scope of the invention. Indeed, numerous  
10 modifications and variations in the practice of the invention are expected to occur to those skilled in the art upon consideration of the present preferred embodiments. Consequently, the only limitations which should be placed upon the scope of the invention are those which appear in the appended claims.

15

All references cited within the body of the instant specification are hereby incorporated by reference in their entirety.

20

25

30